


Projectile Motion

Horizontal Motion

- What would happen if we shot a cannon ball at velocity, v , horizontally in a world without gravity?
 - It would travel horizontally with constant velocity, v




The diagram shows a cannon on a cliff. A red dashed line with an arrow at the end extends horizontally from the cannon, labeled "Inertial or gravity-free path". Below the cliff, a solid black line represents the ground. A red caption at the bottom reads: "In the absence of gravity, an object in motion will continue in motion with the same speed and in the same direction."

Vertical Motion

- What would happen if we dropped a cannon ball from a cliff?
 - It would fall and accelerate with an acceleration, g


Horizontal and Vertical Motion

- What happens when an object is launched horizontally with gravity?
 - The object follows a parabolic path starting from its launching point and eventually ending on the ground



So what does this mean?

- Since gravity only occurs in the vertical direction, it can only affect the vertical motion



- This means that we will treat the horizontal and vertical components of velocity separately

| | Horizontal | Vertical |
|---------------------|------------|---|
| Acceleration | No | Yes g, down (-9.81 ms ⁻²) |
| Velocity | Constant | Changing |

Example

- A cannon ball is launched with a horizontal velocity of 50 m/s from the top of a 10 m high cliff. Determine the distance from the bottom of the cliff where the cannon ball lands.

We will treat this situation as two separate problems: a horizontal one and a vertical one.

- | | |
|--------------------------|--------------------------------|
| • Horizontal | • Vertical |
| • $v_x = 50 \text{ m/s}$ | • $v_y = 0$ |
| • $a = 0$ | • $a = g = -9.8 \text{ m/s}^2$ |
| • $d_x = ?$ | • $d_y = -10 \text{ m}$ |
| • $t = ?$ | • $t = ?$ |

We have enough information to solve for time, t , vertically.

- Vertical

$$d = vt + \frac{1}{2}at^2$$
$$-10 = \frac{1}{2}(-9.8)t^2$$
$$t^2 = 2.04$$
$$t = 1.43 \text{ s}$$

- The time it takes for the object to fall and hit the ground is the same as the horizontal time
 - The object stops moving horizontally once the object has hit the ground
- That means that we can now solve for the horizontal distance

- Horizontal

$$v = \frac{d}{t}$$

$$50 = \frac{d}{1.43}$$

$$d_x = 71.5 \text{ m}$$

- But what if the object is launched at an angle?
 - No problem, we treat it exactly the same way

Example

- A cannon ball is launched with a velocity of 50 m/s at an angle of 30° from the horizontal from the top of a 10 m high cliff. Determine the distance from the bottom of the cliff where the cannon ball lands.

- Once again, we need to separate the horizontal and vertical components
- This time, however, the initial velocity is a vector at an angle
- That means that we have a velocity in both the horizontal and vertical directions

So, let's write down what we know

- | | |
|-------------------------|------------------------------------|
| • Horizontal | • Vertical |
| • $v_x = 50\cos 30$ m/s | • $v_y = 50\sin 30$ m/s |
| • $a = 0$ | • $a = -g = -9.8$ m/s ² |
| • $d_x = ?$ | • $d_y = -10$ m |
| • $t = ?$ | • $t = ?$ |

Once again, we have enough information to solve for time, t , vertically.

- vertical

$$d = vt + \frac{1}{2}at^2$$

$$-10 = (50 \sin 30)t + \frac{1}{2}(-9.8)t^2$$

$$4.905t^2 - 25t - 10 = 0$$

- We have to solve this using the quadratic formula

$$\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$t = \frac{-(-25) \pm \sqrt{(-25)^2 - 4(4.9)(-10)}}{2(4.9)}$$

$$t = \begin{cases} -0.37 \text{ s} \\ 5.47 \text{ s} \end{cases}$$

- Since time cannot be negative, the only value that makes sense is 5.47 s
- Once again, the horizontal part takes the same amount of time
- So now we can solve the horizontal part

- Horizontal

$$v = \frac{d}{t}$$

$$50 \cos 30 = \frac{d}{5.47}$$

$$d_x = 237 \text{ m}$$

- We will use this technique to solve all projectile motion problems
